

STIMUKIL[®] FLY BAIT

ACTIVE INGREDIENTS:

(Z)-9-tricosene.....	0.040%
Methomyl (S-Methyl-N-[(Methylcarbamoyl)oxy]thioacetimidate).....	1.000%

INERT INGREDIENTS	<u>98.960%</u>
TOTAL	100.000%

CAUTION KEEP OUT OF REACH OF CHILDREN

Read Entire Label for Additional Cautions

EPA REGISTRATION NO. 53871-3

EPA ESTABLISHMENT NO. 53871-AZ-01

Manufactured By: TROY BIOSCIENCES INCORPORATED, 113 SOUTH 47TH AVENUE, PHOENIX, AZ 85043

PRECAUTIONARY STATEMENTS HAZARDOUS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing dust. Avoid contact with skin, eyes or clothing. Wash thoroughly after handling and before eating, drinking, and use of tobacco products or using the toilet. **Do not use this product around or inside homes, or any other place where children are likely to be present.**

For commercial use only. Use only in areas inaccessible to food producing animals. Do not contaminate feed and foodstuffs. Do not apply where poultry or animals, especially dogs and young calves, can pick it up or lick it. Wear long sleeved shirt, long pants, shoes, socks, and gloves during loading, application, repairing, and cleaning of application equipment and disposal of the pesticide.

Do not use in edible product areas of food processing plants or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed. **IMPORTANT!** If pesticide comes in contact with skin, wash off with soap and water.

FIRST AID

If in Eyes: Flush eyes with plenty of water for 15 minutes. Get medical attention if irritation persists.

If on Skin: Remove contaminated clothing and wash skin immediately with soap and water. Wash contaminated clothing before reuse.

If Swallowed: Call a physician or poison control center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching the back of the throat with finger. Never induce vomiting or give anything by mouth to an unconscious person.

NOTE TO PHYSICIAN: METHOMYL is a cholinesterase inhibitor. ATROPINE is an antidote.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or by disposal. **Pesticide Disposal:** Pesticide or Rinsate that cannot be used according to label instructions must be disposed of according to applicable Federal, State or local governmental procedures. **Waste Disposal:** Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **Container Disposal:** Triple rinse and dispose of according to local governmental procedures.

Bait Station Disposal: The bait station is reusable. When the container becomes cracked or broken, triple rinse and dispose of according to local government procedures.

ENVIRONMENTAL HAZARDS

Accessible areas treated with this bait are hazardous to birds and other wildlife. Use with care when applying in areas frequented by wildlife or adjacent to any body of water. Collect, cover or incorporate granules spilled on the soil surface. This pesticide is toxic to birds. Do not apply directly to water, or to areas where surface water is present or to intertidal area below the mean high water mark. Do not contaminate water by cleaning of equipment and when disposing of equipment wash-waters. Do not contaminate water by disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. STIMUKIL Fly Bait is an improved fly bait formula containing a sex-attractant and a feeding synergist, which encourages male and female flies to remain in the treated area and stimulates vigorous feeding on the bait. **STIMUKIL Fly Bait should not be used around or inside homes, or any other place where children are likely to be present.** Specific uses are described in more detail below:

BAIT STATION USE: Do not use around or inside homes or other places where children are likely to be present. Pour 25 grams or 2 Tablespoons STIMUKIL Fly Bait into Fly Bait Station. Replace cover carefully to avoid spillage of bait. Use the product at a rate of approximately two bait stations per 500 square feet of fly feeding area. Hang Fly Bait Stations out of reach of food producing animals or pets (at least 4 feet off the ground). Do not place bait station within reach of children. When fly populations are high, the STIMUKIL Fly Bait Station should be replenished with STIMUKIL Fly Bait every 3-5 days or when the bait is no longer visible in the bait station. The dead flies and any remaining bait should be disposed of as recommended above and STIMUKIL Fly Bait added to the Fly Bait Station. Use STIMUKIL Fly Bait Station around the outside of broiler houses, kennels, canneries, beverage processing plants, meat and poultry processing plants, food processing plants and refuse Dumpsters[®] associated with these establishments, and inside of caged layer houses. May also be used in fenced or otherwise enclosed areas inaccessible to children, and around the outside of restaurants, supermarkets, commissaries, and bakeries.

SCATTER BAIT USE: Use of scatter bait is limited solely to commercial establishments where children will not be present, such as feed lots, poultry houses, livestock barns, canneries, beverage plants, meat and poultry processing establishments, commercial refuse dumpsters which are fenced or otherwise enclosed, and food processing plants. Do not use around commercial dumpsters that are not enclosed. Do not use around dumpsters in residential settings. Distribute bait from container or use other device to avoid handling bait directly. Scatter bait (do not apply in piles) at a rate of 1/4 pound per 500 square feet of area. Do not treat areas where animals can feed on bait. Do not apply directly to water. Keep children and pets clear from scatter bait treatment areas. When fly populations are high the bait should be replenished every 3-5 days or when the bait is no longer visible in the treated area.

WARRANTY

Troy Biosciences Incorporated makes no warranty, express or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risks of use and handling of this material when such use and handling are contrary to label instructions.

STIMUKIL IS A REGISTERED TRADEMARK OF TROY BIOSCIENCES INCORPORATED



TROY BIOSCIENCES
INCORPORATED